

# BMJ

## Male subfertility

Anthony Hirsh

BMJ 2003;327:669-672  
doi:10.1136/bmj.327.7416.669

---

Updated information and services can be found at:  
<http://bmj.com/cgi/content/full/327/7416/669>

---

*These include:*

### References

This article cites 2 articles, 1 of which can be accessed free at:  
<http://bmj.com/cgi/content/full/327/7416/669#BIBL>

2 online articles that cite this article can be accessed at:  
<http://bmj.com/cgi/content/full/327/7416/669#otherarticles>

### Rapid responses

4 rapid responses have been posted to this article, which you can access for free at:  
<http://bmj.com/cgi/content/full/327/7416/669#responses>

You can respond to this article at:  
<http://bmj.com/cgi/eletter-submit/327/7416/669>

### Email alerting service

Receive free email alerts when new articles cite this article - sign up in the box at the top left of the article

---

### Topic collections

Articles on similar topics can be found in the following collections

[Other sexual medicine](#) (150 articles)  
[Reproductive medicine](#) (363 articles)

---

### Correction

A correction has been published for this article. The contents of the correction have been appended to the original article in this reprint. The correction is available online at:  
<http://bmj.com/cgi/content/full/327/7424/1165-a>

---

### Notes

---

To order reprints follow the "Request Permissions" link in the navigation box

To subscribe to *BMJ* go to:  
<http://resources.bmj.com/bmj/subscribers>

## ABC of subfertility

### Male subfertility

Anthony Hirsh

Abnormal semen quality or sexual dysfunction are contributing factors in about half of subfertile couples. As natural pregnancy is substantially reduced in these cases, the man should be assessed by an appropriately trained gynaecologist in a reproductive medicine clinic, or by a clinical andrologist. Subfertile men often defer consultations because they perceive subfertility as a threat to their masculinity. Consultations should help them to distinguish between fertility and virility, which may ease their anxiety. However, achievement of a wanted pregnancy is more likely to restore manly feelings.

### Causes of male subfertility

Subfertility affects one in 20 men. Idiopathic oligoasthenoteratozoospermia is the commonest cause of male subfertility. Although sexual function is normal, there is a reduced count of mainly dysfunctional spermatozoa. Reduced fertilising capacity is related to raised concentrations of reactive oxygen species in semen, which may damage the cell membrane. Abnormal sperm morphology—an indicator of deranged sperm production or maturation—is also associated with reduced fertilising capacity. Less common types of male subfertility are caused by testicular or genital tract infection, disease, or abnormalities. Systemic disease, external factors (such as drugs, lifestyle, etc), or combinations of these also result in male subfertility. Male subfertility is rarely caused by endocrine deficiency.


Falling sperm counts have not affected global fertility, although the effect of increased oestrogenic compounds in drinking water is of concern because the incidence of cryptorchidism and testicular cancer is increasing.

#### Semen analysis terminology

- Normozoospermia—All semen parameters normal
- Oligozoospermia—Reduced sperm numbers  
Mild to moderate: 5-20 million/ml of semen  
Severe: <5 million/ml of semen
- Asthenozoospermia—Reduced sperm motility
- Teratozoospermia—Increased abnormal forms of sperm
- Oligoasthenoteratozoospermia—Sperm variables all subnormal
- Azoospermia—No sperm in semen
- Aspermia (anejaculation)—No ejaculate (ejaculation failure)
- Leucocytospermia—Increased white cells in semen
- Necrozoospermia—All sperm are non-viable or non-motile

### Clinical assessment

History taking should include frequency of coitus, erectile function, ejaculation, scrotal disorders or surgery, urinary symptoms, past illnesses, lifestyle factors, and any drugs taken. Physical examination should seek signs of hypogonadism (small testes), hypoandrogenism (lack of facial or body hair), systemic disease, and abnormalities of the penis or testicles. Testicular sizes are assessed by length (cm) or volume (ml) and are measured with an orchidometer. Because of the risk of cancer, a urological opinion is essential if there is an intratesticular lump or the testes are undescended or absent. Scrotal ultrasonography is helpful in confirming a varicocele or



Sperm morphology is related to the fertilising capacity by in vitro fertilisation. (A=normal sperm head; B=abnormal head; C=globozoospermia—a rare syndrome in which all sperm heads lack acrosome caps and cannot fertilise)

**Semen analysis is the cornerstone of male fertility assessment and is often the trigger to refer patients for a specialist opinion. Semen samples are best sent to laboratories linked with infertility services. Ideally two samples around six weeks apart (unless the first is unequivocally normal) are required. The samples should be produced by masturbation after three days' abstinence. Men may use non-spermicidal condoms if they have difficulty with, or religious objections to, masturbation**

#### Normal seminal fluid analysis (World Health Organization, 2002)

- Volume > 2 ml
- Sperm concentration > 20 million/ml
- Sperm motility > 50% progressive or > 25% rapidly progressive
- Morphology (strict criteria) > 15% normal forms
- White blood cells < 1 million/ml
- Immunobead or mixed antiglobulin reaction test\* < 10% coated

\*Tests for the presence of antibodies coating the sperm



A simple orchidometer is a 4 cm long (20 ml) ovoid used to assess testis size in subfertile men. Normal testes are more than 4 cm and firm in consistency. Abnormal testes are soft and smaller. Both testes should be carefully examined to exclude a tumour

## Clinical review

testicular tumour. Transrectal ultrasonography of the prostate may identify the cause of a low volume ejaculate. Serum follicle stimulating hormone is a useful index of impaired spermatogenesis. Genetic screening (karyotype, or DNA analysis for Y chromosome microdeletions or cystic fibrosis) is indicated for men with severe oligozoospermia and most men with azoospermia.

## Treatment options for subfertile men

As all couples hope and prefer to conceive naturally, a specific diagnosis should be sought and corrected where appropriate. However, a couple with subfertility of longer than three years, or with a non-reversible form of subfertility is unlikely to conceive spontaneously and should join an assisted conception programme without delay.

### What doesn't work

- Abstaining from coitus until ovulation does not improve the semen or likelihood of conception. Increasing coital frequency (alternate days) supplies more viable spermatozoa that normally remain motile in the female tract for two to three days
- Treatment with gonadotrophin injections, androgens (mesterolone) or antiestrogens (clomifene or tamoxifen) is not indicated because although they may improve the sperm count, fertility rates are not improved as the spermatozoa remain dysfunctional

### Stopping adverse drugs and drug misuse

Several drugs impair spermatogenesis or sexual function. Most common are sulfasalazine and anabolic steroids when misused by athletes. These effects are reversible, allowing fertility to return to normal in six to 12 months if the drugs are withdrawn. Chemotherapy and radiotherapy damage spermatogenesis, hence sperm banking should be offered to male patients with cancer irrespective of sperm quality.

### Timing and lifestyle changes

Most cases of mild to moderate oligozoospermia are idiopathic, but transient oligozoospermia can follow influenza or a major illness and improves within three to six months. The incidence of spontaneous conception each month is 1-2% and justifies conservative or empirical treatment for younger couples. The incidence also explains the powerful placebo effect of some treatments. Advice can be given on lifestyle changes and on the avoidance of fertility impairing drugs.

### Treating accessory gland infection

With the increased prevalence of chlamydia, accessory gland infection may cause partial obstruction, focal epididymitis, or subclinical prostatitis. Semen cultures are rarely useful but antibiotic treatment (for example, doxycycline, erythromycin, ciprofloxacin) is often given empirically. Antioxidants (for example, vitamins C and E) absorb reactive oxygen species and are purported to improve sperm motility, although no convincing evidence exists that pregnancy rates are improved.


### Assisted conception

Assisted conception gives most infertile men the chance of biological fatherhood, and it is most successful if the woman is under 35 years. The method indicated depends on the quantity and quality of sperm isolated from the semen after "washing" or density gradient techniques. The resulting sperm preparations have improved counts of morphologically normal progressively motile spermatozoa.

### Genetics and male infertility

Clinical diagnosis	Genetic tests	Most common defects	Incidence (%)
Congenital bilateral absence of vas deferens (CBAVD)	Cystic fibrosis (CFTR gene)	ΔF508, R117H	66
Non-obstructive azoospermia	Karyotype	47, XXY AZFa, AZFb*, AZFc	15-30 10-15
Severe (<5M/ml) oligozoospermia	Karyotype  Y chromosome microdeletions	47, XXY Translocation Partial AZFb, AZFc	1-2 0.2-0.4 7-10

CFTR = cystic fibrosis transmembrane conductance regulator; AZF = azoospermia factor; \*AZFb the most severe (DAZ gene—deleted in azoospermia) causes the most severe defects of spermatogenesis; AZFc causes the mildest defects of spermatogenesis



Autosomal Robertsonian reciprocal translocation may be associated with poor sperm quality and subfertility

### Drugs that can impair male fertility

- Impaired spermatogenesis—Sulfasalazine, methotrexate, nitrofurantoin, colchicine, chemotherapy
- Pituitary suppression—Testosterone injections, gonadotrophin releasing hormone analogues
- Antiandrogenic effects—Cimetidine, spironolactone
- Ejaculation failure— $\alpha$  blockers, antidepressants, phenothiazines
- Erectile dysfunction— $\beta$  blockers, thiazide diuretics, metoclopramide
- Drugs of misuse—Anabolic steroids, cannabis, heroin, cocaine

### Conservative measures for men with suboptimal semen analyses

- Stop smoking—Nicotine reduces seminal plasma antioxidants
- Have frequent intercourse—Increases output of non-senile spermatozoa
- Reduce alcohol intake—Alcohol can suppress spermatogenesis
- Wear boxer shorts and avoid hot baths—Heat suppresses spermatogenesis
- Avoid pesticides, herbicides, heat, and radiation at work—All impair spermatogenesis

### *Intrauterine insemination*

In mild or moderate oligozoospermia some spermatozoa are functionally normal. Intrauterine insemination is feasible with preparations of three to five million progressively motile sperm. The woman must have at least one normal patent fallopian tube for successful interuterine insemination. Combined with ovarian stimulation, three to four cycles of intrauterine insemination result in conception in 15-30% of couples.

### *In vitro fertilisation and intracytoplasmic sperm injection*

A prepared sample containing around one to two million motile sperm is required for adequate oocyte fertilisation with in vitro fertilisation. Not surprisingly, fertilisation is lowest when it is the man who is infertile. However, intracytoplasmic sperm injection needs only one viable sperm for microinjection into each egg. The technique is indicated if the semen preparation yields too few normal motile sperm for in vitro fertilisation, as occurs in severe oligozoospermia. Intracytoplasmic sperm injection is appropriate after unexpectedly, poor, or absent fertilisation in vitro. It is also an effective technique for men with azoospermia by using spermatozoa that have been surgically retrieved from the epididymis or testis.

## Severe oligoasthenoteratozoospermia

Most cases of severe oligoasthenoteratozoospermia are caused by idiopathic testicular abnormality or disorder. Genetic tests are abnormal in 7-10% of men with sperm counts less than 5 million/ml. Severely impaired sperm motility may be caused by antibodies, chronic prostatitis, or rare recessive intrinsic defects of the sperm tail linked to sinopulmonary disease (for example, Kartagener's syndrome or Young's syndrome). Treatment of severe oligoasthenoteratozoospermia rarely improves the semen quality, but intracytoplasmic sperm injection is often successful even if only few weakly motile spermatozoa can be isolated from the ejaculate.

## Azoospermia

Azoospermia (absence of sperm from the semen) can be caused by hypothalamic-pituitary failure, testicular failure, or testicular obstruction. Testicular size and concentration of follicle stimulating hormone determine the clinical diagnosis. Although azoospermia is uncommon, 75% of men who have the condition now have the opportunity of biological fatherhood through assisted conception techniques.

### **Hypogonadotrophic hypogonadism**

Hypogonadotrophic hypogonadism is rare but can be treated with gonadotrophin injections or by administering gonadotrophin releasing hormone by infusion pump. Natural conceptions often occur within a year of treatment because any spermatozoa secreted will be functionally normal.

### **Obstructive azoospermia**


Men with obstructive azoospermia have normal spermatogenesis and hence normal size testes, normal concentrations of serum follicle stimulating hormone, and they are normally virilised. If neither vas is palpable, congenital bilateral absence of the vas deferens is diagnosed, which cannot be corrected surgically. As two thirds of men with palpable congenital bilateral absence of the vas deferens carry cystic fibrosis mutations, both partners require screening. Other cases of obstructive azoospermia occur after vasectomy or they are caused by epididymal obstruction after chlamydia or gonorrhoea. Vasectomy reversal will return sperm to the

**Testing for sperm antibodies is controversial. False positive results occur and antibodies do not necessarily impair sperm function. Antibodies may be found in genital infections and obstructions, but specific treatment is of limited value. Corticosteroids have been used successfully for high titre sperm antibodies. However, severe side effects may occur, including bilateral necrosis of the hip and gastric ulceration. In vitro fertilisation is a more effective and safer way to achieve pregnancy**

**Live birth rates of 20-30% per in vitro fertilisation cycle are achievable, even in severe oligozoospermia or azoospermia, if the woman is under 35 years**

### **Varicocele controversy**

Varicoceles occur in 15% of men in general and in 30% of subfertile men. Varicoceles probably impair spermatogenesis by increasing the temperature in the scrotum. Achievement of pregnancies is often attributed to varicocele surgery because semen quality may improve after surgery and because conceptions occur in 15% of infertile men with or without surgery. However, clinical trials are equivocal. Varicoceles can be ligated or embolised, but as it may take one to two years for the couple to achieve a pregnancy, clinicians may recommend assisted conception if the woman is older than 35 years



Microsurgical vasovasectomy (vasectomy reversal). Using an operating microscope, the cut and often fibrosed ends of the vas deferens are dissected free from surrounding tissue and anastomosed using fine nylon sutures to re-establish patency. The small squares on the graph paper are 1 mm wide

ejaculate in 80-90% of men, and pregnancies occur in 40-50% of couples in one to two years. Testicular exploration may be indicated for other obstructions because reconstructive surgery results in sperm positive semen in 30-50% of cases, and pregnancies in 20-25% of couples. If necessary, sperm retrieved from the epididymis during these reconstructive procedures can be frozen for future intracytoplasmic sperm injection cycles.

### Non-obstructive azoospermia

Non-obstructive azoospermia may be caused by cryptorchidism, Klinefelter's syndrome (47,XXY), or Y chromosome deletions after chemotherapy or radiotherapy—for example, for lymphoma or testicular cancer. However, many cases of non-obstructive azoospermia are idiopathic. Multiple testicular biopsy may show scattered foci of spermatogenesis in about half the cases with potential for surgical sperm retrieval and intracytoplasmic sperm injection. Men with non-obstructive azoospermia should have genetic testing as 15-30% of them have sex chromosome aneuploidy or Y chromosome deletions.

### Surgical sperm retrieval

Surgical sperm retrieval for intracytoplasmic sperm injection is indicated in obstructive azoospermia where spermatogenesis is usually normal, or non-obstructive azoospermia where spermatogenesis is present on biopsy in 30-50% of cases. Results of intracytoplasmic sperm injection using surgically retrieved sperm are similar to cycles where ejaculated sperm is used. Percutaneous epididymal or testicular sperm aspiration or extraction are usually feasible under local anaesthetic and sedation.

## Sexual dysfunction


Most men with erectile or ejaculation failure have normal sperm function and are managed according to whether ejaculation can be stimulated. Men unable to produce the sample for in vitro fertilisation usually respond to sildenafil.

In retrograde ejaculation, where the emission enters the bladder because of non-surgical sphincter failure (for example, in diabetes), oral sympathomimetics (for example, pseudoephedrine) may close the incompetent bladder neck and produce antegrade ejaculation. Retrograde ejaculation caused by anatomical sphincter defects (for example, after prostatectomy or other bladder neck incision) is managed by intrauterine insemination. The sperm that are used are isolated from post-ejaculation urine, which is suitably alkalinised by oral sodium bicarbonate and adjusted for osmolarity.

In men whose spinal cord is injured, semen is usually obtained with a vibrator when vaginal insemination at home may be successful. If this fails, and in cases of aspermia caused by pelvic injury or multiple sclerosis, rectal electrostimulation usually provides semen suitable for assisted conception. If ejaculation is not induced, sperm can be retrieved by vas deferens aspiration or by testicular aspiration for intracytoplasmic sperm injection.

## Donor insemination

Donor insemination is the principal choice for the 1 in 200 infertile men (and their partners) who have no sperm because of traumatic or congenital anorchia or total germ cell aplasia. As donor semen is selected for high quality, the live birth rate of 7-15% per cycle is largely dependent on the fertility of the woman. Adoption is the only other option at present.



Testicular biopsy of non-obstructive azoospermia showing Sertoli cell only syndrome with a focus of spermatogenesis. All biopsies obtained for testicular sperm extraction in non-obstructive azoospermia are assessed histologically because of the increased prevalence (0.4-1.1%) of carcinoma in situ in subfertile men

### Further reading

- Hargreave TB, Mills JA. Investigating and managing infertility in general practice. *BMJ* 1998;316:1438-41
- Hirsh AV. Investigation and therapeutic options for infertile men presenting in assisted conception units. In: Brinsden PR, ed. *In-vitro fertilisation and assisted reproduction*. 2nd ed. London: Parthenon, 1999
- Hull MGR, Glazener CMA, Kelly NJ, Conway DI, Foster PA, Hinton RA, et al. Population study of causes, treatment, and outcome of infertility. *BMJ* 1985;291:1693-7
- Royal College of Obstetricians and Gynaecologists. *Evidence-based guidelines: initial investigation and management of the infertile couple*. London: RCOG, 1998
- Royal College of Obstetricians and Gynaecologists. *Evidence-based guidelines: management of infertility in secondary care*. London: RCOG, 1998
- Royal College of Obstetricians and Gynaecologists. *Evidence-based guidelines: management of infertility in tertiary care*. London: RCOG, 2000
- Rowe PJ, Comhaire FH, Hargreave TB, Mahmoud AMA. *WHO manual for the standard investigation, diagnosis and management of the infertile male*. Cambridge: Cambridge University Press, 2000
- Skakkebaek NE, Giwercman A, de Kretser D. Pathogenesis and management of male infertility. *Lancet* 1994;343:1473-9
- Vale J, Hirsh AV. *Male sexual dysfunction*. Oxford: Blackwell Science, 2001

The scanning electron micrographs of sperm morphology are published with permission of Alpha from *Scientists in Reproductive Medicine* newsletters 1996 and 2000. The testicular biopsy showing non-obstructive azoospermia is at a magnification x 400 and is reprinted with permission from Dr K Shah, London.

Anthony Hirsh is consultant to the andrology clinic at Whipps Cross Hospital, London, and honorary senior lecturer at King's, Guy's, and St Thomas's School of Medicine, London.

The ABC of subfertility is edited by Peter Braude, professor and head of department of women's health, Guy's, King's, and St Thomas's School of Medicine, London, and Alison Taylor, consultant in reproductive medicine and director of the Guy's and St Thomas's assisted conception unit. The series will be published as a book in the winter.

Competing interests: None declared.

*BMJ* 2003;327:669-72

## Summary points

Methods of communicating health risks to health policy makers have been neglected

Decision makers require easily understandable measures that show the impact of risk factors for disease on populations to help guide the allocation of resources according to local health needs

The population impact number of eliminating a risk factor (PIN-ER-*t*) is “the potential number of disease events prevented in your population over the next *t* years by eliminating a risk factor”

The PIN-ER-*t* can be used to show the impact of a range of risk factors in different populations and to compare the potential benefits of individual and population approaches to prevention

We have reported that individual clinicians are not as influenced by the presentation of risk in population terms as they are by relative risk (Heller et al, submitted for publication), while others have found that the “number needed to treat” statistic (which also relies on measures of absolute risk) is poorly understood by doctors and lay people.<sup>29 30</sup> It remains for us to examine whether new measures of population impact like PIN-ER-*t* can be more easily understood and used in health policy related decision making than traditional methods of communicating risk. We are developing a research programme to explore this further.

**Contributors and sources:** The authors work at the Evidence for Population Health Unit, aiming to develop a public health counterpart to evidence based medicine. The measure described here is one of a series of population impact measures developed to use evidence combined with routinely collected data to provide local context to measures of risk and benefit and support public health policy decision making.

**Competing interests:** None declared.

- Edwards A, Elwyn G, Mulley A. Explaining risks: turning numerical data into meaningful pictures. *BMJ* 2002;324:827-30.
- Fahey T, Griffiths S, Peters TJ. Evidence based purchasing: understanding results of clinical trials and systematic reviews. *BMJ* 1995;311:1056-9.
- Last JM. *A dictionary of epidemiology*. Oxford: Oxford University Press, 2001.
- McPherson K, Britton A, Casner L. *Coronary heart disease. Estimating the impact of changes in risk factors*. London: Stationery Office, National Heart Forum, 2002.
- Schoenbach VJ. Relating risk factors to health. *Epidemiolog.net* 2002. [www.epidemiolog.net/evolving/RelatingRiskFactorstoHealth.pdf](http://www.epidemiolog.net/evolving/RelatingRiskFactorstoHealth.pdf) (last updated 23 Jun 2003).
- Gail MH, Benichou J. *Encyclopaedia of epidemiologic methods*. Chichester: John Wiley, 2000.
- Cook R, Sackett D. The number needed to treat: a clinically useful measure of treatment effect. *BMJ* 1995;310:452-4.
- Milward L, Kelly M, Nutbeam D. *Public health intervention research: the evidence*. London: Health Development Agency, 2001.
- Levin ML. The occurrence of lung cancer in man. *Acta Unio Int Contra Cancrum* 1953;19:531.
- Walter SD. Choice of effect measures for epidemiological data. *J Clin Epidemiol* 2000;52:931-9.
- Miettinen OS. Proportion of disease caused or prevented by a given exposure, trait or intervention. *Am J Epidemiol* 1974;99:325-32.
- Armitage P, Berry G, Matthews JNS. *Statistical methods in medical research*. Oxford: Blackwell Science, 2002.
- Heller RF, Dobson AJ, Atia J, Page JH. Impact numbers: measures of risk factor impact on the whole population from case control and cohort studies. *J Epidemiol Community Health* 2002;56:606-10.
- Department of Health. *Compendium of clinical and health indicators 2001*. London: DoH, 2002. (<http://www.nchod.nhs.uk>)

- Heller RF, McElduff P, Edwards R. Impact of upward social mobility on population mortality: analysis with routine data. *BMJ* 2002;325:134.
- Walker A, O'Brien M, Traynor J, Fox K, Goddard E, Foster K. *Living in Britain 2001: health survey for England 2001*. London: Stationery Office, Office for National Statistics, 2002.
- Marang-van de Mheen PJ, Davey-Smith G, Hart CL. The health impact of smoking in manual and non-manual social class men and women: a test of the Blaxter hypothesis. *Soc Sci Med* 1999;48:1851-6.
- Department of Health. *Health survey for England 1998: cardiovascular disease*. London: Stationery Office, 1999.
- Kannel WB, Neaton JD, Wentworth D, Thomas HE, Stamler J, Hulley SB, et al. Overall coronary heart disease mortality rates in relation to major risk factors in 325,348 men screened for MRFT. *Am Heart J* 1986;112:825-36.
- Chen Z, Peto R, Collins R, MacMahon S, Lu J, Li W. Serum cholesterol concentration and coronary heart disease in populations with low cholesterol concentrations. *BMJ* 1991;303:276-82.
- Rose G. Sick individuals and sick populations. *Int J Epidemiol* 1985;14:32-8.
- Murray CJ, Lauer JA, Hutubessy RCW, Niessen L, Tomijima N, Rogers A, et al. Effectiveness and costs of interventions to lower systolic blood pressure and cholesterol: a global and regional analysis on reduction of cardiovascular-disease risk. *Lancet* 2003;361:717-25.
- Ezzati M, Lopez AD. Measuring the accumulated hazards of smoking: global and regional estimates for 2000. *Tobacco Control* 2003;12:79-85.
- Peto R, Lopez AD, Boreham J, Thun M, Heath C Jr. Mortality from tobacco in developed countries: indirect estimation from national vital statistics. *Lancet* 1992;339:1268-78.
- Walter SD. Prevention for multifactorial diseases. *Am J Epidemiol* 1980;112:409-16.
- Morgenstern H, Bursic E. A method for using epidemiologic data to estimate the potential impact of an intervention on the health status of a target population. *J Community Health* 1982;7:292-309.
- Heller RF, Page JH. A population perspective to evidence based medicine: “evidence for population health.” *J Epidemiol Community Health* 2002;56:45-7.
- Heller RF, Edwards R, McElduff P. Implementing guidelines in primary care: can population impact measures help? *BMC Public Health* 2003;3:7. <http://www.biomedcentral.com/1471-2458/3/7>
- Nexoe J, Gyrd-Hansen D, Kragstrup J, Kristiansen IS, Nielsen JB. Danish GPs' perception of disease risk and benefit of prevention. *Fam Pract* 2002;19:3-6.
- Kristiansen IS, Gyrd-Hansen D, Nexoe J, Nielsen JB. Number needed to treat: easily understood and intuitively meaningful? Theoretical considerations and a randomised trial. *J Clin Epidemiol* 2002;55:888-92. (Accepted 19 August 2003)

## Corrections and clarifications

*Parathyroid hormone alone is as effective as combination in treating osteoporosis*

We enthusiastically added a reference to this news article by Scott Gottlieb to help readers locate the study being reported (27 September, p 700). Unfortunately, although we got the year and volume of the *New England Journal of Medicine* right, we published the wrong page numbers. The correct reference is 2003;349:1207-15.

*ABC of subfertility: male subfertility*

Two errors crept into in this article by Anthony Hirsh (20 September, pp 669-72). Firstly, we incorrectly inserted an extra word in the caption to the figure on page 670; the caption should read: “Autosomal Robertsonian translocations may be associated with poor sperm quality and subfertility.” Secondly, we made a dog's dinner of the caption to the figure on page 671. The photograph in fact shows a “microsurgical vasovasostomy for vasectomy reversal.”

*General practitioners and occupational health professionals*

We inadvertently typed the word “health” instead of “medicine” when we inserted the competing interests for one of the authors of this editorial by Jeremy Beach and David Watt (9 August, pp 302-3). Professor Beach is in fact an assistant editor of the journal *Occupational Medicine*.